

Name \_\_\_\_\_  
Literacy Lab #13 - Dinosaur Emissions

Date \_\_\_\_\_  
Earth Science - Breed - 2012/2013

**Directions:** Take a few minutes to read the article below either online (or on the back of this page.) Write responses to the statements or questions below. Cut/copy/paste is not allowed – use your own words and thoughts, based in research if needed.

Read more: <http://www.sciencedaily.com/releases/2012/05/120507102324.htm>

**Fact-finding:** List three facts that you learned in this article.

1.

2.

3.

**Vocabulary:** List and define three unfamiliar words in the space below.

**Implications:** What are your feelings about this “discovery”? Express your feelings (tactfully) about whether this is an advancement of science or a bad idea.

# Gaseous Emissions from Dinosaurs May Have Warmed Prehistoric Earth

*ScienceDaily* (May 7, 2012) — Sauropod dinosaurs could in principle have produced enough of the greenhouse gas methane to warm the climate many millions of years ago, at a time when Earth was warm and wet. That's according to calculations reported in the May 8th issue of *Current Biology*, a Cell Press publication.

The hulking sauropods, distinctive for their enormous size and unusually long necks, were widespread about 150 million years ago. As in cows, methane-producing microbes aided the sauropods' digestion by fermenting their plant food.

"A simple mathematical model suggests that the microbes living in sauropod dinosaurs may have produced enough methane to have an important effect on the Mesozoic climate," said Dave Wilkinson of Liverpool John Moores University. "Indeed, our calculations suggest that these dinosaurs could have produced more methane than all modern sources -- both natural and man-made -- put together."

Wilkinson and study coauthor Graeme Ruxton from the University of St Andrews were studying sauropod ecology when a question dawned on them: If modern cows produce enough methane gas to be of interest to climate scientists, what about sauropods? They teamed up with methane expert Euan Nisbet at the University of London to work out the numbers.

"Clearly, trying to estimate this for animals that are unlike anything living has to be a bit of an educated guess," Wilkinson said.

Animal physiologists have studied methane production from a range of modern animals to derive equations that predict methane production from animals of different sizes. It turns out that those calculations depend only on the total mass of the animals in question. A medium-sized sauropod weighed something like 20,000 kilograms, and sauropods lived in densities ranging from a few large adults to a few tens of individuals per square kilometer.

Wilkinson, Ruxton, and Nisbet therefore calculate global methane emissions from sauropods to have been 520 million tons (520 Tg) per year, comparable to total modern methane emissions. Before industry took off on modern Earth about 150 years ago,

methane emissions were roughly 200 Tg per year. By comparison, modern ruminant animals, including cows, goats, giraffes, and others, produce methane emission of 50 to 100 Tg per year.

The study's conclusions not only show "just how strange and wonderful the workings of the planet are" but also serve as a useful reminder for the importance of microbes and methane for global climate, the researchers say.

Journal reference: Wilkinson et al.: "Could methane produced by sauropod dinosaurs have helped drive Mesozoic climate warmth?"

*Share this story on **Facebook**, **Twitter**, and **Google**:*

*Other social bookmarking and sharing tools:*

#### **Story Source:**

The above story is reprinted from [materials](#) provided by [Cell Press](#), via [EurekAlert!](#), a service of AAAS.

*Note: Materials may be edited for content and length. For further information, please contact the source cited above.*

*Note: If no author is given, the source is cited instead.*

**Disclaimer:** *Views expressed in this article do not necessarily reflect those of ScienceDaily or its staff.*